

DFPD/93/TH/69

hept-th/9403079

**THE HIGGS MODEL FOR ANYONS AND LIOUVILLE ACTION:
CHAOTIC SPECTRUM, ENERGY GAP AND EXCLUSION PRINCIPLE[¶]**

MARCO MATONE*

Department of Physics “G. Galilei” - Istituto Nazionale di Fisica Nucleare

University of Padova

Via Marzolo, 8 - 35131 Padova, Italy

ABSTRACT

Geodesic completeness and self-adjointness imply that the Hamiltonian for anyons is the Laplacian with respect to the Weil-Petersson metric. This metric is complete on the Deligne-Mumford compactification of moduli (configuration) space. The structure of this compactification fixes the possible anyon configurations. This allows us to identify anyons with singularities (elliptic points with ramification q^{-1}) in the Poincaré metric implying that anyon spectrum is chaotic for $n \geq 3$. Furthermore, the bound on the holomorphic sectional curvature of moduli spaces implies a gap in the energy spectrum. For $q = 0$ (punctures) anyons are infinitely separated in the Poincaré metric (hard-core). This indicates that the exclusion principle has a geometrical interpretation. Finally we give the differential equation satisfied by the generating function for volumes of the configuration space of anyons.

[¶]Partly Supported by the European Community Research Programme *Gauge Theories, applied supersymmetry and quantum gravity*, contract SC1-CT92-0789

*e-mail: matone@padova.infn.it, mvxpd5::matone

1. Let us begin with the following remark

a. *The configuration space of n anyons is the space of n unordered points in $\hat{\mathbf{C}} = \mathbf{C} \cup \{\infty\}$*

$$M_n = (\hat{\mathbf{C}}^n \setminus \Delta_n) / \text{Symm}(n),$$

with Δ_n the diagonal subset where two or more punctures coincide.

b. *The Liouville action on the Riemann sphere with n -punctures evaluated on the classical solution is the Kähler potential for the natural metric (the Weil-Petersson metric) on¹*

$$\mathcal{M}_n = (\hat{\mathbf{C}}^n \setminus \Delta_n) / \text{Symm}(n) \times PSL(2, \mathbf{C}).$$

This remark implies that starting from anyons on $\hat{\mathbf{C}}$ one can recover the two-form associated to the natural metric on the configuration space by first computing the Poincaré metric e^φ on the punctured sphere and then, after evaluating the Liouville action for $\phi = \varphi$, computing the curvature two-form of the Hermitian line bundle on $\mathcal{M}_n$ defined by the classical action (see below).

To understand the physical relevance of this remark we notice that the quantum Hamiltonian for n anyons is proportional to the covariant Laplacian on M_n . In the case of anyons on the thrice punctured Riemann sphere we have

$$H = -\frac{1}{2m} \Delta_{WP}, \tag{1}$$

where Δ_{WP} is the Weil-Petersson Laplacian.

A crucial point is the physical requirement of self-adjointness of the Hamiltonian. In our approach this requirement is satisfied by considering the Hamiltonian defined on $\overline{\mathcal{M}}_n$ where $\partial\overline{\mathcal{M}}_n = \overline{\mathcal{M}}_n \setminus \mathcal{M}_n$ denotes the Deligne-Mumford compactification of $\mathcal{M}_n$. In particular we recall that the Weil-Petersson metric defines a Hermitian inner product and has a completion to the boundary $\partial\overline{\mathcal{M}}_n$. Thus (1) describes a well-defined physical problem. In particular the structure of the Deligne-Mumford compactification fixes possible anyon-configurations. We notice that in the degeneration limit punctures never collide (hard-core). This is a consequence of the requirement of stability. On the other hand this can be seen as a consequence of the fact that the Liouville equation has delta-singularities at the punctures (hard-core).

¹The $PSL(2, \mathbf{C})$ group reflects the Möbius symmetry of the Riemann sphere. Thus one can consider $\mathcal{M}_n$ as the configuration space for $n - 3$ anyons on the Riemann sphere with 3-punctures.

The connection between anyons and Liouville arises also in considering the critically coupled Abelian Higgs model in (2+1)-dimensions where the space M_n plays a crucial role in the analysis of the n^{th} topological sector of the theory. Remarkably $f = 2\text{Re} \log \phi$, with ϕ the Higgs field, satisfies the modified (non covariant) Liouville equation

$$f_{z\bar{z}} = e^f - 1, \quad (2)$$

which is the second Bogomol'nyi equation. Notice that the non covariance (the -1 in (2)) is due to the Higgs mass. In [1] Samols has studied the metric on vortex moduli space in the Abelian Higgs model. In [2] Manton has proposed a model for anyons which is based on the structure of such a space. The Lagrangian density of the Abelian Higgs model is

$$\mathcal{L} = \frac{1}{2} D_\mu \phi \overline{D^\mu \phi} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{8} (|\phi|^2 - 1)^2, \quad \phi = \phi_1 + i\phi_2, \quad (3)$$

where $D_\mu \phi = (\partial_\mu - iA_\mu)\phi$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, and the metric has signature $(1, -1, -1)$. In the temporal gauge the finiteness of the energy implies that at infinity the Higgs field is a pure phase. The magnetic flux through $\mathbf{R}^2$ is $\int F_{12} = 2\pi n$ where n is the winding number labelling the topological sectors of the map $|\phi| : S_\infty^1 \rightarrow U(1)$. In the static configuration $\dot{A}_i = 0$, $\dot{\phi} = 0$, the energy has the lower bound $E \geq \pi|n|$. The critical case $E = \pi|n|$ arises when the Bogomol'nyi equations

$$(D_1 + \text{sgn}(n)iD_2)\phi = 0, \quad F_{12} + \text{sgn}(n)\frac{1}{2}(|\phi|^2 - 1) = 0, \quad (4)$$

are satisfied. It is crucial that in the n^{th} topological sector the space of smooth solutions is a manifold $\widetilde{M}_n$ of complex dimension n . In particular, each solution is uniquely specified by the n unordered points $\{z_k\}$ where the Higgs field is zero [3]. The same happens in considering the Liouville equation for the Poincaré metric on Riemann surfaces. In particular, due to the uniqueness of the solution of the Poincaré metric, to each complex structure of the n -punctured Riemann sphere (the unordered set of n points) corresponds a solution of the Liouville equation (see below for details).

Topologically $\widetilde{M}_n$ coincides with $\widehat{\mathbf{C}}^n$. It is a remarkable fact that the Abelian Higgs model (almost) provides a smooth metric and $U(1)$ gauge field on $\widetilde{M}_n$. In particular the kinetic energy induces the metric

$$ds^2 = \frac{1}{2} \int_{\mathbf{C}} (d\phi_a d\phi_a + dA_i dA_i).$$

The field evolution is described by geodesic motion on $\widetilde{M}_n$. Notice that this corresponds to deformation of the complex structure of punctured spheres. The z_k 's are good coordinates

only on the subspace M_n . Good global coordinates on M_n are provided by the coefficients of the polynomial [4] $P_n(z) = \sum_{k=0}^n w_k z^k \equiv \prod_{k=1}^n (z - z_k)$. We stress that the choice of good coordinates is a subtle point. Actually, as we will see, instead of working with this kind of structure it is much more convenient to work with the Weil-Petersson metric. This is directly related to Liouville theory and Fuchsian groups. This structure is at the origin of chaotic spectra for anyons.

In [1] Samols introduced the metric $ds^2 = \frac{1}{2} \sum_{r,s=1}^n \left(\delta_{rs} + 2 \frac{\partial \bar{b}_r}{\partial z_s} \right) dz_r d\bar{z}_s$ on $\widetilde{M}_n$ where the b_r 's satisfy the equations

$$\frac{\partial b_r}{\partial \bar{z}_s} = \frac{\partial \bar{b}_s}{\partial z_r}. \quad (5)$$

2. Let us denote by H the upper half-plane and with Γ a finitely generated Fuchsian group. A Riemann surface isomorphic to the quotient H/Γ has the Poincaré metric $\hat{g}$ as the unique metric with scalar curvature $R_{\hat{g}} = -1$ compatible with its complex structure. This implies the uniqueness of the solution of the Liouville equation on Σ

$$\partial_z \partial_{\bar{z}} \varphi(z, \bar{z}) = \frac{1}{2} e^{\varphi(z, \bar{z})}. \quad (6)$$

Let us consider the punctured Riemann sphere $\Sigma = \widehat{\mathbf{C}} \setminus \{z_1, \dots, z_n\}$, $\widehat{\mathbf{C}} \equiv \mathbf{C} \cup \{\infty\}$. The moduli space of n -punctured spheres is the space of classes of isomorphic Σ 's, that is

$$\mathcal{M}_n = \{(z_1, \dots, z_n) \in \widehat{\mathbf{C}}^n | z_j \neq z_k \text{ for } j \neq k\} / \text{Symm}(n) \times PSL(2, \mathbf{C}), \quad (7)$$

where $\text{Symm}(n)$ acts by permuting $\{z_1, \dots, z_n\}$ whereas $PSL(2, \mathbf{C})$ acts by linear fractional transformations. By $PSL(2, \mathbf{C})$ we can recover the 'standard normalization': $z_{n-2} = 0$, $z_{n-1} = 1$ and $z_n = \infty$. For the classical Liouville tensor $T^F = \varphi_{zz} - \varphi_z^2/2$ we have

$$T^F(z) = \sum_{k=1}^{n-1} \left(\frac{1}{2(z - z_k)^2} + \frac{c_k}{z - z_k} \right), \quad \lim_{z \rightarrow \infty} T^F(z) = \frac{1}{2z^2} + \frac{c_n}{z^3} + \mathcal{O}\left(\frac{1}{|z|^4}\right).$$

The c_k 's, called accessory parameters, are functions on the space

$$V^{(n)} = \{(z_1, \dots, z_{n-3}) \in \mathbf{C}^{n-3} | z_j \neq 0, 1; z_j \neq z_k, \text{ for } j \neq k\}, \quad (8)$$

which is a covering of $\mathcal{M}_n$

$$\mathcal{M}_n \cong V^{(n)} / \text{Symm}(n), \quad (9)$$

where the action of $\text{Symm}(n)$ on $V^{(n)}$ is defined by comparing (7) with (9).

The Liouville action on the Riemann spheres with n -punctures

$$S^{(n)} = \lim_{r \rightarrow 0} \left[\int_{\Sigma_r} \left(\partial_z \phi \partial_{\bar{z}} \phi + e^\phi \right) + 2\pi(n \log r + 2(n-2) \log |\log r|) \right],$$

$\Sigma_r = \Sigma \setminus \left(\bigcup_{i=1}^{n-1} \{z \mid |z - z_i| < r\} \cup \{z \mid |z| > r^{-1}\} \right)$ has the following property [5]

$$-\frac{1}{2\pi} \frac{\partial S_{cl}^{(n)}}{\partial z_k} = c_k, \quad k = 1, \dots, n-3, \quad (10)$$

where the c_k 's are the accessory parameters. Since $S_{cl}^{(n)}$ is real, eq.(10) yields $\frac{\partial c_j}{\partial \bar{z}_k} = \frac{\partial \bar{c}_k}{\partial z_j}$ corresponding to the Samols equations (5). It turns out that [5]

$$\frac{\partial c_j}{\partial \bar{z}_k} = \frac{1}{2\pi} \left\langle \frac{\partial}{\partial z_j}, \frac{\partial}{\partial z_k} \right\rangle^{(n)}, \quad j, k = 1, \dots, n-3, \quad (11)$$

where the brackets denote the Weil-Petersson metric on the Teichmüller space T_n projected onto $V^{(n)}$. Therefore by (10)

$$\omega_{WP}^{(n)} = \frac{i}{2} \bar{\partial} \partial S_{cl}^{(n)} = -i\pi \sum_{j,k=1}^{n-3} \frac{\partial c_k}{\partial \bar{z}_j} d\bar{z}_j \wedge dz_k, \quad (12)$$

where ω_{WP} is the Weil-Petersson two-form on $V^{(n)}$.

3. The symplectic structure above allows us to consider the Poisson bracket relations [7]

$$\{c_j, c_k\}_{WP} = 0, \quad \{c_j, z_k\}_{WP} = \frac{i}{\pi} \delta_{jk}, \quad (13)$$

where the brackets are defined with respect to ω_{WP} . These relations suggest performing the geometric quantization of space $\mathcal{M}_n$. To do this we must define a suitable line bundle. Let $\sigma_{k,n}$ be the element of $Symm(n)$ interchanging k and n . Let us consider the function [6]

$$f_{\sigma_{k,n}} = \frac{\prod_{j \neq k, j=1}^{n-3} (z_j - z_k)^2}{(z_k(z_k - 1))^{n-4}}, \quad k = 1, \dots, n-3, \quad f_{\sigma_{n-2,n}} = \prod_{j=1}^{n-3} z_j^2, \quad f_{\sigma_{n-1,n}} = \prod_{j=1}^{n-3} (z_j - 1)^2. \quad (14)$$

Extension by the composition $f_{\sigma_1 \sigma_2} = (f_{\sigma_1} \circ \sigma_2) f_{\sigma_2}$ defines a 1-cocycle $\{f_\sigma\}_{\sigma \in Symm(n)}$ of $Symm(n)$ [6]. Let us consider the holomorphic line bundle $\mathcal{L}_n = V^{(n)} \times \mathbf{C} / Symm(n)$, on $\mathcal{M}_n$ where the action of $\sigma \in Symm(n)$ is defined by $(x, z) \rightarrow (\sigma x, f_\sigma(x)z)$, $x \in V^{(n)}$, $z \in \mathbf{C}$. Since $\exp(S_{cl}^{(n)} \circ \sigma / \pi) |f_\sigma|^2 = \exp(S_{cl}^{(n)} / \pi)$ it follows that $\exp(S_{cl}^{(n)} / \pi)$ is a Hermitian metric

in the line bundle $\mathcal{L}_n \rightarrow \mathcal{M}_n$. By (10) $\exp(S_{cl}^{(n)}/\pi)$ has connection form $-2\sum_i c_i dz_i$ and, by (12), curvature two-form $-\frac{2i}{\pi}\omega_{WP}$ [6]. The covariant derivatives are

$$\mathcal{D}_k = \partial_{z_k} - \partial_{z_k} \frac{S_{cl}^{(n)}}{\pi} = \partial_{z_k} + 2c_k, \quad \overline{\mathcal{D}}_k = \partial_{\bar{z}_k}. \quad (15)$$

In the geometric quantization the Hilbert states are sections of $\mathcal{L}_n$ annihilated by ‘half’ of the derivatives (polarization). The natural choice is

$$\mathcal{H} = \left\{ \psi \in \mathcal{L}_n \mid \overline{\mathcal{D}}_k \psi = 0 \right\}, \quad (16)$$

with inner product

$$\langle \psi_1 | \psi_2 \rangle = \frac{1}{n!(n-3)!} \int_{\mathcal{M}_n} d(WP) e^{-\frac{S_{cl}^{(n)}}{\pi}} \overline{\psi}_1 \psi_2, \quad (17)$$

where

$$d(WP) \equiv \left(\bigwedge^{n-3} \frac{i}{2} \overline{\partial} \partial S_{cl}^{(n)} \right), \quad (18)$$

that by (12) is the Weil-Petersson volume form. An alternative to (18) is to use the volume form $\bigwedge^{n-3} \omega$ where

$$\omega = \frac{i}{2} \overline{\partial} \partial \log \det \left\| \frac{\partial^2 S_{cl}^{(n)}}{\partial \bar{z}_j \partial z_k} \right\|. \quad (19)$$

However, in general, the correspondence principle can be proved only if the scalar product is defined with respect to the volume form associated with the Kähler potential [8]. This means that in our case we must use the Weil-Petersson volume form.

We conclude the discussion on the geometric quantization of $\mathcal{M}_n$ by noticing that an interesting alternative for the polarization choice in the geometric quantization above is

$$\mathcal{H} = \left\{ \tilde{\psi} \in \mathcal{L}_n \mid \mathcal{D}_k \tilde{\psi} = 0 \right\}. \quad (20)$$

In this case the states and the classical Liouville action are related by

$$\partial_{z_k} S_{cl}^{(n)} = \pi \partial_{z_k} \log \tilde{\psi}. \quad (21)$$

4. A crucial aspect that should be further investigated concerns the classical-quantum interplay arising both in Liouville and anyon theories². In [9, 10] it has been emphasized that

²Vortices are unlabelled particles not only quantum mechanically but also classically [2].

the regularization arising at the classical level for the Liouville action is strictly related to the conformal properties both of quantum and classical (Poincaré metric) Liouville operators.

The approach considered here makes it possible to define the conformal weight associated to anyons [10]

$$\Delta(q) = \frac{1 - q^2}{2h}, \quad (22)$$

where q^{-1} is the ramification index of the anyon (elliptic point). For $q = 0$ the point is parabolic (puncture). In this case the geodesic distance between an arbitrary point and an anyon is infinity in the natural (Poincaré) metric. In other words, if z_k is the coordinate of the anyon (with index $q = 0$) then

$$\lim_{Q \rightarrow z_k} d(P, Q) = \infty. \quad (23)$$

In particular two anyons with $q = 0$ are infinitely separated! This means that for spin $1/2h$, particles cannot coalesce in the Poincaré metric (hard-core). This connection between spin and hard-core resembles the Pauli exclusion principle. Actually this hard-core is encoded in our approach. The reason is that the Liouville equation on the sphere with the punctures filled-in has additional terms given by δ -functions at the punctures. These aspects should be useful in investigating the standard approach to anyon [11].

5. We now consider the energy spectrum of anyons. As we have seen Liouville theory arises in the framework of negatively curved surfaces. This suggests to consider the Gutzwiller model [12] for chaos. To see this notice that in the case of one anyon moving on the thrice-punctured sphere the Hamiltonian (1) reduces to the Laplacian on the Riemann sphere with punctures fixed at 0, 1 and ∞ . Note that fixed punctures may be considered as infinitely massive anyons. The crucial point is that in this case the Hamiltonian corresponds to the Laplacian on a Riemann surface of negative curvature. Therefore we can apply the results in [12] concerning quantum chaos on Riemann surfaces.

This result can be extended to the general situation of multi-anyon spectra. The point is that moduli space of higher genus (i.e. $h \geq 2$) is negatively curved. In particular the holomorphic sectional curvatures are bounded away from zero by a negative constant [13]. Therefore it seems that the previous result about chaoticity for anyons is a general situation which holds for $n \geq 3$.

The previous bound on the holomorphic sectional curvature suggests that there is a mass-gap in the anyon spectra. This is a direct consequence of the infrared properties of negatively

curved surfaces. Actually this is the case if one considers one anyon moving on the thrice punctured sphere. In particular it is well-known that the eigenvalues of the Laplacian have a gap in the spectrum.

Let us make further remarks on the anyon dynamics. The point is that it depends on the geodesic (complex) structure. This structure is determined by the position of the anyons themselves, resulting structure is fascinating and highly nontrivial. Actually we can consider anyons as singularities in the metric with a self gravitational interaction.

Results from the theory of moduli spaces allow to compute the generating function for the volumes of configuration space of anyons (factorized by the trivial infinity coming from the $PSL(2, \mathbf{C})$ symmetry). In particular it turns out that the generating function satisfies the differential equation [14]

$$g'' = \frac{g'^2 t - g g' + g' t}{t(t - g)}, \quad (24)$$

where

$$g(t) = \sum_{k=3}^{\infty} \frac{k(k-2)}{(k-3)!} t^{k-1} \int_{\mathcal{M}_k} \left(\frac{i \bar{\partial} \partial S_{cl}^{(k)}}{2\pi^2} \right)^{k-3}, \quad (25)$$

where “ $\int_{\mathcal{M}_3} 1$ ” $\equiv \frac{1}{6}$.

Another interesting aspect is the connection between Liouville and Higgs. This provides a way to consider the Higgs model in a 2D gravity framework.

Finally we stress that the relation between Liouville theory and anyons arises also in a different context (see [15] and references therein).

Acknowledgements. I would like to thank C. Grosche for remarks on chaos, R. Jackiw for bringing ref.[15] to my attention and P.A. Marchetti for stimulating discussions.

References

- [1] T.M. Samols, Phys. Lett. **244B** (1990) 285; Comm. Math. Phys. **145** (1992) 149; *Soliton Scattering*, Cambridge University PhD Thesis, 1991.
- [2] N.S. Manton, *Vortices And Anyons*, preprint DAMTP 91-19.
- [3] A. Jaffe and C.H. Taubes, *Vortices And Monopoles*, Boston, Birkhäuser, 1980.
C.H. Taubes, Comm. Math. Phys. **72** (1980) 277; **75** (1980) 207.

- [4] N.J. Hitchin, in ‘Proceedings of the Lefschetz Centennial Conference’. Part I. Cont. Math. vol. **58** AMS 1986.
- [5] P.G. Zograf and L.A. Takhtajan, Math. USSR Sbornik, **60** (1988) 143.
- [6] P.G. Zograf, Leningrad Math. J. **1** (1990) 941.
- [7] L.A. Takhtajan, Proc. Symp. Pure Math. AMS **49**, part 1 (1989) 581.
- [8] F.A. Berezin, Math. USSR Izvestia **8** (1974) 1109.
- [9] M. Matone, *Uniformization Theory And 2D Gravity. I. Liouville Action And Intersection Numbers*, preprint IC-Math/8-92, DFPD/92/TH/41, hep-th/9306150.
- [10] M. Matone, *Quantum Riemann Surfaces, 2D Gravity And The Geometrical Origin Of Minimal Models*, preprint DFPD/93/TH/62, hep-th/9309096.
- [11] J. Fröhlich and P.A. Marchetti, Lett. Math. Phys. **16** (1988) 347; Comm. Math. Phys. **121** (1989) 177.
- [12] M.C. Gutzwiller, *Chaos In Classical And Quantum Mechanics*, Springer-Verlag 1990. N. Balazs and A. Voros, Phys. Rep. **143** (1986) 109. J. Bolte and C. Grosche, *Selberg Trace Formula For Bordered Riemann Surfaces: Hyperbolic, Elliptic And Parabolic Conjugacy Classes, And Determinants Of Maass-Laplacians*, Desy 92-118, SISSA/139/92/FM, to appear in Comm. Math. Phys.
- [13] H. Royden, Lect. Notes Math. **185** (1971) 125. S. Wolpert, Invent. Math. **85** (1986) 119.
- [14] M. Matone, *Nonperturbative Model Of Liouville Gravity*, DFPD/93/TH/72, hep-th/9402081.
- [15] R. Jackiw, Int. J. Mod. Phys. **B5** (1991) 1617.